

Multivariable Calculus Quiz 5

Due Thursday 10/22 12:00 p.m.

Please show all work to receive full credit.

Don't forget that the timer on Gradescope includes time needed to scan and submit your work!

Problem 1 (3 pts)

Suppose the temperature on a surface is given by the function

$$T(x, y) = 100e^{-(x^2+y^2)}.$$

Now suppose there is a bug that takes the path of a curve given by the function

$$\vec{r}(t) = \langle t \cos(2t), t \sin(2t) \rangle.$$

What is the rate at which the temperature changes as the bug moves?

Problem 2 (2 pts)

In fluid dynamics, *laminar flow* is characterized by fluid particles following smooth paths in layers, in contrast to something like turbulent flow. Considering laminar flow in a cylinder, the *resistance* R to the flow is related to the length and radius of the cylinder by the *Hagen–Poiseuille law*. More specifically, the resistance is given by the following function:

$$R(w, r) = k \frac{w}{r^4}$$

where w is the length of the cylinder, r is the radius of the cylinder, and k is a constant.

Give the linearization of R at the point (w_0, r_0) .

Problem 3 (5 pts)

Let $\rho(x, y, z, t)$ be a smooth function (that is, all its partial derivatives exist and are differentiable). Let

$$\tilde{\nabla} \rho = \left\langle \frac{\partial \rho}{\partial x}, \frac{\partial \rho}{\partial y}, \frac{\partial \rho}{\partial z} \right\rangle$$

denote the gradient of ρ **only in the first three variables** (this is not a standard notation, I made this up myself).

Let $\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle$ be a smooth curve, and let $\mathbf{v}(t) = \mathbf{r}'(t)$ denote the velocity of the curve. You can think of $\rho(x, y, z, t)$ as a function that describes the density of a fluid at the point (x, y, z) in space, and at time t . While the curve $\mathbf{r}(t)$ is the path of a point particle in the flow.

For the following questions we restrict to studying the change of ρ on the path $\mathbf{r}(t)$ (otherwise none of the expressions in the questions will make sense).

1. Use the chain rule to show that

$$\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \mathbf{v} \cdot \tilde{\nabla} \rho.$$

2. We say that the fluid is *incompressible* if

$$\frac{d\rho}{dt} = 0.$$

- Suppose that the density function ρ is independent of the spatial variables x, y, z , i.e. $\rho(x, y, z, t) = \rho(t)$. Use the result of Part 1. to give the condition on ρ for the fluid to be incompressible.
- Now suppose instead that the density function ρ is **only dependent on the spatial variables** x, y, z , i.e. $\rho(x, y, z, t) = \rho(x, y, z)$. In this case if a flow is incompressible, we call it a *stratified flow*. Use the result of Part 1. to give the condition for ρ and \mathbf{v} for a stratified flow. Note: Say something about the relation between ρ and \mathbf{v} .